


Fourier series in $L^2(S^1)$

$\mathcal{B} = (e_n)_{n \in \mathbb{Z}}$ for $e_n: x \mapsto e^{2\pi i n x} \in L^2(S^1)$

$$S^1 = \mathbb{R}/\mathbb{Z}$$

Q Why is $e_n \in L^2(S^1)$?

Q Why is $\mathcal{B}$ an orthonormal system?



$$\int_0^1 |e_n|^2 = \int_0^1 |e_n^2| = \int_0^1 |e_{2n}| = \int_0^1 1 = 1 < \infty$$

• $\|e_n\| = 1$ by $\nearrow$

$$\langle e_n, e_m \rangle = \int_0^1 e_n \overline{e_m} = \int_0^1 e_{n-m} = \dots = 0$$

For $f \in L^2(S')$, the n -th Fourier coefficient of f is

$$\hat{f}(n) = \langle f, e_n \rangle$$

$$= \int_0^1 f(x) \overline{e_n(x)} \, dx$$

$$= \int_0^1 f(x) e^{-2\pi i n x} \, dx \in \mathbb{C}$$

The N -th Fourier polynomial of f is the projection of f onto $\text{span}_{\mathbb{C}}\{e_{-N}, \dots, e_N\}$, i.e.

$$f_N(x) := \sum_{n=-N}^N \hat{f}(n) e_n(x).$$

The Fourier series of f is

$$F[f](x) := \lim_{N \rightarrow \infty} f_N(x) = \sum_{n \in \mathbb{Z}} \hat{f}(n) e_n(x).$$

Facts • Call $\text{span}_{\mathbb{C}} \{e_{-N}, \dots, e_N\}$ the space of trigonometric polynomials of degree N . For $W \leq V$, proj_W^V is the vector in W closest to v : $\text{proj}_W^V = \underset{w \in W}{\text{argmin}} \|v - w\|$. inner prod space

$$\|f_N\|^2 = \sum_{n=-N}^N |\hat{f}(n)|^2$$

monotonically
increasing seq

Thus $f_N(x)$ is the trig poly of degree N best approximating f in L^2 norm. (Best Approx'n Thm)

Bessel's inequality $\|f_N\| \leq \|f\|$ ~~IOU~~.

- By the proof of our main theorem from Friday,
 $\sum_{n \in \mathbb{Z}} |\hat{f}(n)|^2$ converges so, in particular, we have the
Riemann-Lebesgue lemma $\lim_{n \rightarrow \pm \infty} \hat{f}(n) = 0$.

- If $\mathcal{B}$ is an orthonormal basis, also get
Parseval's identity : $\|f\|^2 = \sum_{n \in \mathbb{Z}} |\hat{f}(n)|^2$.

Forthcoming goal:

Inversion Thm $\mathcal{B}$ is an orthonormal basis for $L^2(S')$.

Cor $f = \mathcal{F}[f]$ in $L^2(S')$, and $L^2(S') \cong \ell^2(\mathbb{Z})$
 $f \mapsto (\hat{f}(n))_{n \in \mathbb{Z}}$.



Equality in L^2 is not pointwise convergence.

Later we will show $F[f]$ converges absolutely & uniformly to f for $f \in C^1(S^1)$ — and something similar holds for $C^0(S^1)$!

not an algebraic kernel


Convolutions & kernels

Fourier series have a useful relationship with an operation called convolution:

Fourier series: conv'n $::$ rings: mult'n

Defn For $f, g \in C^0(S^1)$, the convolution $f * g \in C^0(S^1)$

is given by $(f * g)(x) := \int_0^1 f(x-t)g(t) dt$.



Moral ex $t \mapsto f(x-t)g(t) \in C^0(S^1)$, $f * g \in C^0(S^1)$.

Thm • Convolution is linear in each variable

• Convolution is commutative $f * g = g * f$

• Convolution is associative $(f * g) * h = f * (g * h)$

• For $f \in C^1(S^1)$, $g \in C^0(S^1)$, have $f * g \in C^1(S^1)$ and

$$(f * g)' = f' * g.$$

• For $f, g \in C^0(S^1)$, $\widehat{f * g}(n) = \hat{f}(n) \hat{g}(n)$.

IF HW. $\square$

WELCOME TO
WISHFUL
THINKING

Suppose we have a Dirac delta function δ such that for $f \in C^0(S^1)$,

$$\int_{-1/2}^{1/2} \delta(x) f(x) dx = f(0).$$

$$\text{Then } f * \delta(x) = \int_{-1/2}^{1/2} f(x-t) \delta(t) dt = f(x-0) = f(x)$$
$$\Rightarrow \hat{f}(n) \hat{\delta}(n) = \hat{f}(n) \Rightarrow \hat{\delta}(n) = 1$$

i.e. $f * \delta = f$ — δ is an identity for the convolution op'n.

Now suppose we want to show $\lim_{n \rightarrow \infty} f_n = f$, and we have K_n such that $f * K_n = f_n$ and $\lim_{n \rightarrow \infty} K_n = \delta$. Then

$$\lim_{n \rightarrow \infty} f_n = \lim_{n \rightarrow \infty} f * K_n = f * \lim_{n \rightarrow \infty} K_n = f * \delta = f.$$


Defn A Dirac kernel on S' is a sequence of continuous

$K_n : [-1/2, 1/2] \rightarrow \mathbb{R}$ such that

(1) $K_n \geq 0$

(2) $\int_{-1/2}^{1/2} K_n(x) dx = 1.$

(3) For any $\delta > 0$, $\lim_{n \rightarrow \infty} \int_{\delta \leq |x| \leq 1/2} K_n(x) dx = 0$



Q What do we learn about $\int_{-\delta}^{\delta} K_n(x) dx$? $\xrightarrow[\substack{\delta > 0 \\ n \rightarrow \infty}]{}$ 1

Defn The Dirichlet kernel $\{D_N \mid N \in \mathbb{N}\}$ is

$$D_N(x) := \sum_{n=-N}^N e_n(x).$$

The Fajér kernel $\{F_N \mid N \geq 1\}$ is

$$F_N(x) = \frac{1}{N} \sum_{k=0}^{N-1} D_k(x) \quad (\text{see demo})$$

2025. II. 5

Thm For $f \in C^0(S^1)$, $f * D_N = f_N$ and

$$f * F_N = \frac{1}{N} \sum_{k=0}^{N-1} f_k.$$

Pf By linearity of convolution, it suffices to show that